

TRIALS FOR PREPARATION OF INACTIVATED EGG DROP SYNDROME VIRUS VACCINE ADAPTED ON SPECIFIC PATHOGEN FREE EMBRYONATED CHICKEN EGGS (SPF-ECE)

NADIA M. IBRAHIM, EMAN A. HASSAN AND ABO-ZIED A. ABO-ZIED

Veterinary Serum and Vaccine Research Institute, ARC, Ministry of Agriculture, Dokki-Giza

(Manuscript received 13 June 2010)

Abstract

Egg drop syndrome – 76 (EDS-76) virus was propagated for three passages on embryonated duck eggs followed by thirty passages on C.E.F cell culture, then, for ten passages on SPF embryonated chicken eggs. The titer and HA activity of the 10th passage was 10^{9.5}/ml and 2¹⁰/ml successively. Inactivation of passage number 10 of propagated virus on SPF-ECE with formaline was adjuvanted with paraffin oil. The prepared vaccine was compared with inactivated oil emulsion EDS76 prepared on embryonated duck eggs for sterility, safety and potency. The potency test was performed by measuring the cellular and humoral immune response. The efficiency of the prepared vaccine was estimated up to six months.

INTRODUCTION

EDS virus has become a major cause of lost egg production throughout the world. It is caused by an adenovirus. The disease is characterized by the production of thin-shelled or shell-less eggs (Van Eck *et al.*, 1976).

The virus appeared to be transmitted vertically through the egg. The virus often remained latent until birds were approaching peak egg production (Calnek, 1978).

An oil adjuvant inactivated vaccine is widely used to giving good protection against clinical EDS (Christensen, 1998).

The birds were vaccinated between 14 and 16 weeks of age (Lee and Hopkins, 1982). HI antibody response can be detected by the 7th day after vaccination with peak titers between the 2nd and 5th weeks. The vaccinal immunity lasts at least one year (Baxendale *et al.*, 1980, Khalaf *et al.*, 1982).

The aim of the present study is directed to the preparation and evaluation of inactivated EDS vaccine on SPF embryonated chicken eggs.

MATERIALS AND METHODS

1. **Chicks:** two hundred susceptible 21-days old Hubbard chicks were used for vaccine evaluation.

2. Virus strain: EDS-76 virus strain was supplied by the Central Veterinary Laboratory, Weybridge, England.

3. Embryos

- a. Embryonated chicken specific pathogen free (SPF) eggs were obtained from Ministry of Agriculture, Koum Osheim, Fayoum, Egypt). Nine to 10 days old SPF-ECE were used for preparing chicken embryo fibroblast cell culture according to Plowright and Ferris (1959) and modified by Osman *et al.* (1985) and also used for propagation of EDS-76 virus.
- b. Embryonated duck eggs: These were obtained from United Company for Poultry Production and used for propagation and preparation of inactivated oil EDS-76 vaccine.

4. Cell culture media, reagents and solutions

- a. Minimum Essential Medium (MEM): it was used as growth medium with 10% newborn calf serum and as maintenance medium with 2-3% newborn calf serum in pH 7.2. It was supplied by Sigma.
- b. Hank's balanced salt solution (HBSS): it was used for virus titration and was prepared according to Hank and Wallace (1949).
- c. Trypsin (1:250): it was used in primary cell culture preparation at a concentration of 0.25% according to Lennette (1964).

Methods

1. Adaptation and propagation of EDS-76 on chicken embryo fibroblast was carried out for thirty passages, and then, adapted and propagated to thirty and fifteen passages on SPF-ECE for 10 passages, and titration to each passage on SPF-ECE to detect HA activity to each passage was applied (Wo, 1995).
2. Preparation of different types of vaccines
 - a. Inactivated oil emulsion EDS-76 on SPF-ECE.
 - b. Inactivated oil emulsion EDS-76 on embryonated duck egg was applied according to Rozhest Vensk (1984).
3. Vaccination of chicks
 - a. Each chick was inoculated with 0.5 ml of the prepared inactivated vaccine by s/c route .
 - b. Safety test
Each chick was inoculated with double dose (1 ml) of the prepared inactivated vaccine by s/c route.

The previous vaccines were tested for sterility according to OIE and for immunological effect with serum neutralization test according to (Rossiter *et al.*, 1985), and for Haemagglutination inhibition test HI according to (Anon, 1971), lymphocyte blastogenesis assay (Lee, 1994), protection percent and keeping quality (Lee and Hopkins, 1982).

RESULTS

Table 1. Propagation and titration of EDS-76 virus in SPF embryonated chicken eggs.

No. of passages	Log ₁₀ EID ₅₀ /ml	HA (log ₂)
1	3.50	3
2	4.10	3
3	4.10	5
4	6.10	6
5	8.50	7
6	8.50	8
7	7.80	9
8	9.00	10
9	9.30	10
10	9.50	10

Table 2 . Sterility of the prepared EDS vaccine.

Media	Inactivated oil emulsion	
	Embryonated SPF propagated vaccine	Embryonated duck eggs vaccine
Nutrient agar media,	NC	NC
Thioglycolate broth	NT	NT
Sabaraoud's glucose agar	NC	NC
Grey media	NC	NC

NC = no colonies

NT = no turbidity

Table 3 . Lymphocyte blastogenesis of chicks vaccinated with the prepared vaccines.

Group	Type of vaccines used	Weeks post-vaccination		
		1 st	2 nd	3 rd
1	Inactivated oil emulsion EDS-76 propagated on SPF embryonated chicken eggs	0.72325	0.93325	0.96125
2	Inactivated oil emulsion EDS-76 on embryonated duck eggs	0.57320	0.84520	0.72340
3	Control non-vaccinated	0.18550	0.19200	0.18150

Table 6. Rate of protection for the prepared EDS-76 virus vaccines.

G	type of used vaccines	1 st month			2 nd month			3 rd month			4 th month		
		No. of challenged chicks	Survived	Protection %	No. of challenged chicks	Survived	Protection %	No. of challenged	Survived	Protection %	No. of challenged	Survived	Protection %
1	Inactivated oil emulsion EDS-76 on SPF embryonated chicken eggs	5	5	100	5	5	100	5	5	100	5	5	100
	Inactivated oil emulsion EDS-76 on duck eggs	5	5	100	5	5	100	5	5	100	5	5	100
3	Control non-vaccinated	5	0	0	0	0	0	5	0	0	5	0	0

Table 7. Keeping quality of prepared EDS-76 virus vaccines (months).

Group	Type of vaccine used	Temp of storage	Protection %							
			1 st month		2 nd month		3 rd month		4 th month	
			Survived chickens	Protection %	Survived chickens	Protection %	Survived chickens	Protection %	Survived chickens	Protection %
1	Inactivated oil emulsion EDS-76 on SPF embryonated chicken eggs	4 °C	5/5	100	5/5	100	5/5	100	5/5	100
2	Inactivated oil emulsion EDS-76 on duck eggs	4 °C	5/5	100	5/5	100	5/5	100	5/5	100
3	Control	-	0/2	0	0/2	0	0/2	0	0/2	0

DISCUSSION

The aim of the present study was directed to prepare a safe and protective inactivated SPF embryonated chicken egg vaccine against EDS-76, and comparing it with inactivated oil EDS prepared on duck embryonated eggs. The scheme used for preparing the vaccine included the propagation of the adapted EDS-76 virus on CEF cells for 10 passages on SPF-ECE.

Table 1 showed that the titer of the propagated EDS virus on SPF-ECE gradually increased the 1st passage $10^{3.5}$ and reached to maximum in the 10th passage $10^{9.5}$. These results are in agreement with those obtained by Wo (1995).

Table 2 showed that both vaccines were tested for sterility and proved to be free from any contaminants. A high value of lymphocyte blastogenesis for three weeks post-vaccination was shown with the prepared vaccines when compared with control (Table 3). These results were in agreement with Umesh-Kumar *et al.* (1989).

Table 4 showed the evaluation of the humoral immune response HI of chicken vaccinated with the prepared vaccine when compared with vaccine prepared on duck embryonated eggs. It was clear that high titer value of haemagglutination inhibition titer appeared in vaccinated groups for 24th weeks. It was shown that HI titer in chicken vaccinated with ECE vaccine had increased gradually and reached its maximum 2^{11} in 10, 11, 12, weeks and still being high 2^8 to the 24th week, while, HI value of vaccinated chickens with embryonated duck egg vaccine had reached its maximum 2^{10} in 4th -7th till reached 2^7 in 24th week. These results were in agreement with Chen Wen Xian *et al.* (1996).

Table 5 showed that there was negligible difference of neutralizing antibody titre in chicken vaccinated with different vaccines, while, the control group was zero. This result agreed with (Kaur *et al.*, 1997).

Table 6 showed that chicks vaccinated with prepared vaccines gave high protection percentage reaching 100% when vaccinated chicks were challenged with virulent EDS virus after 3 weeks, while, controls did not show any protection. This result agreed with Cook (1983).

By discussing the keeping quality of the prepared inactivated EDS virus vaccines when stored at 4 ° C (Table 7), it was clear that the vaccine was stable and potent for a period of 6 months as protection reached 100%. This result agreed with Khalaf (1981) and Picault *et al.* (1982) who reported that the antigen released over a long period with high immune response after one dose.

We could conclude that the 10th passage of EDS virus on SPF-ECE and the preparation of inactivated oil emulsion vaccine gave safe, potency and immunogenicity. So, from the previous results, we can conclude that the inactivated oil emulsion vaccine prepared on embryonated chicken egg is safe, potent and immunogenic. So, we could use the SPF-ECE instead of embryonated duck eggs.

REFERENCES

1. Anon 1971. Studies of an agent causing mortality among ducklings immune to duck hepatitis virus. *Avian Dis.*, 13: 834-846.
2. Baxendale, W.D. , R. Hein Luhicken and I. Mcpherson. 1980. The results of field trials conducted with an inactivated vaccine against the egg drop syndrome-76 (EDS-76). *Avian Pathol.*, 9: 77-91.
3. Baxendale, W. 1978. Egg drop syndrome-76. *Vet. Rec.*, 102: 285-286.
4. Calnek, B.W. 1978. Haemagglutination inhibition antibodies against an adenovirus (virus-127) in white Pekin ducks in the united states. *Av. Dis.*, 22:798-801.
5. Chen Wenxian, Fanya Fang, Chen Junxia. 1996. Serological measurement of egg drop syndrome-1976 in hens in Jiaquan, Gansu. *Chinese J.Vet.Sci.Tech.*, 26(3): 22-23.
6. Christensen, W.H. 1998. Trial of an inactivated vaccine against egg drop syndrome 76 in New- Zealand. *Vet.J.*, 6: 237-238.
7. Cook, J.K.A. 1983. Egg drop syndrome. 1976. (EDS-76) virus infection in inadequately vaccinated chickens. *Avian Pathol.*, 12(1): 9-16.
8. Hank, J.H. and R.E. Wallace. 1949. Relation of oxygen and temperature in preservation of tissues by refrigeration. *Proc.Soc.Exptl.Bio.Med.*, 71: 196-200.
9. Kaur, A., M.S. Oberoi and Singh Amarjit. 1997. Neutralizing antibody and challenge response to live and inactivated avian adenovirus in broilers. *Trop. Anim. Hlth. Prod.*, 29 (3): 141-146.
10. Khalaf, S.E. 1981. Field and laboratory experiments on immunizing of hens against egg drop syndrome. 1976. Inaugural Dissertation, Tierärztliche Hochschule, Hannover, pp. 89.
11. Khalaf, S. Ed., E.F. Kaleta and O. Steymann. 1982. Comparative studies on the kinetics of haemagglutination inhibition and virus neutralizing antibodies following vaccination of chickens against egg drop syndrome 1976 (EDS-76). *Dev.Biol.Stand.*, 51: 127-137.
12. Lee, A.M.T. and I.G. Hopkins. 1982. Development of a potency test for inactivated egg drop syndrome-76 vaccines. *Develop.Biol.Stand.*, 51: 65-74.
13. Lennette, E.H. 1964. Diagnostic procedure for viral and rickettsial disease. 3rd ed., A Public Health Assoc. Inc. Broadway.
14. Osman, O.A., M.A. Mouaz, S. Athnasiaus and S. Abd El-Ghaffar. 1985. Comparative study of in vitro and in vivo titration of pooled batches of tissue culture rinderpest vaccine. *Al-Azhar J.Pharm.Sci.*, 4: 87-93.

15. Picault, J.P., M. Guihet and G. Benne Jean. 1982. Experiments on the duration of protection conferred by vaccines against aviadenovirus (EDS-76) infection. *Develop.Biol.Stand.*, 51: 139-149.
16. Plowright, W. and R.D. Ferris. 1959. Studies on rinderpest virus in tissue culture. I-Growth and cytopathogenicity. *J.Comp.Pathol.*, 29: 152.
17. Rossiter, P.B., D.M. Tessett and W.P. Taylor. 1985. Microneutralization system for use with different strains of peste des petits ruminants virus and rinder pest virus. *Trop. Anim. Hlth.Prod.*, 17(2): 75-81.
18. Rozhdest, I.K. Venskii. 1984. Inactivating the aviadenovirus of the egg drop syndrome (strain EDS-76). *Veterinaryis Moscow, USSR*, 4: 61-62.
19. Umesh Kumar, K., S. Kishnaswamy and T. Venkata Reddy. 1989. Cell mediated immune response to egg drop syndrome 76 (EDS-76) virus infection in chickens. *Current Science Indian*, 58(8): 431-433.
20. Van Eck, J.H.H., F.G. Davelear, Van den T.A.M. Heplesman, N. Vankol and B.F.H.M.G. Kouwenhoven. 1976. Dropped egg production, soft shelled and shell less eggs associated with appearance of precipitins to a virus in flocks of laying fowl. *Avian Pathol.*, 5: 261-272.
21. Wo. 1995. Methods for the cultivation of infectious laryngeotrachitis virus and egg drop syndrome virus.

محاولة لتحضير لقاح مثبط ضد فيروس مرض تدني البيض في الدواجن على أجنة البيض الخالي من المسببات المرضية

نادية محمد إبراهيم ، إيمان أحمد حسن ، أبو زيد أحمد أبو زيد

معهد بحوث الأمصال واللقاحات البيطرية بالعباسية- مركز البحوث الزراعية- وزارة الزراعة - النقي-جيزة

تم محاولة تمرير فيروس مرض تدني البيض المستضعف على خلايا أجنة الكتاكيت، عشر تمريرات على أجنة البيض الخالي من المسببات المرضية، وقد تم قياس القوة العيارية وقوة التلازن الدموي بعد تمريرة على حدة ، وقد تم تحضير لقاح ميت من التمريرة العاشرة باستخدام الفورمالين وتم حقنه في الكتاكيت القابلة للعدوى وقياس المناعة الخلوية واستخدمت التجارب السيرولوجية (إختبار التعادل المصلي وإختبار منع التلازن الدموي) لقياس المناعة لللقاحات المختبرة على الكتاكيت (لقاح زيتي ميت على أجنة بيض الدجاج ولقاح زيتي ميت على أجنة بيض البط). كما تم إختبار القدرة المناعية للدجاج المحصن باستخدام تجربة التحدي بالفيروس الضاري، وقد تم حفظ اللقاح الميت في درجة 4 م° لمدة ستة أشهر متتالية وإختبار قوة الصد للكتاكيت المحصنة بكل منها .

و توصي الدراسة السابقة بأنه يمكن استخدام أجنة البيض الخالي من المسببات المرضية في تمرير فيروس مرض تدني البيض في الدواجن بدلا" من استخدام أجنة بيض السبط كما توصي بتحضير لقاح مثبط على أجنة البيض الخالي من المسببات المرضية حيث أثبت اللقاح المحضر أنه آمن ، ذو كفاءة عالية و يعطي رد فعل مناعي يكفي لحماية الكتاكيت المحصنة مدة لا تقل عن ستة أشهر .