

EFFICIENCY OF CERTAIN PLANT EXTRACTS AGAINST *CERATITIS CAPITATA* (WIED.) AND *BACTROCERA ZONATA* (SAUNDERS) (DIPTERA: TEPHRITIDAE)

HANAFY, A. H.¹, ANAS A. AHMED¹, A. M. Z. MOSALLAM²
AND A. M. IBRAHIM¹

1. Plant Protection Research Institute, El-Sabahia, Alex.
2. Plant Prot. Res. Inst., ARC, Dokki, Giza

(Manuscript received 12 August 2007)

Abstract

Extracts of six wild plants of *Ambrosia maritima*, *Allium sativum*, *Mentha spicata*, *Myoporum pictum*, *Nerium oleander* and *Rosa gallica* were examined as contact toxic substances against 3-day old pupae and 2-day old adults of both *Ceratitis capitata* and *Bactrocera zonata*. Extracts of *R. gallica* leaves and *A. sativum* bulbils were the most efficient against pupae of *C. capitata* and *B. zonata*, respectively. On the other hand, leaves extract of *N. oleander* was the most effective against adults of both *C. capitata* and *B. zonata* recording the highest cumulative mortality (after 72 hours) of 55.93 and 90.00%, consecutively. As a general trend, adults of *B. zonata* were more susceptible to the tested extracts than that of *C. capitata*.

Key words: Plant extracts- *Ceratitis capitata*-*Bactrocera zonata*- Plant extracts.

INTRODUCTION

Fruit flies such as *C. capitata* and *B. zonata* cause serious damage in fruits of several hosts especially citrus, apricot, peach, mango and guava causing severe reduction in both quantity and quality of infested fruits in Egypt and in many countries.

Generally, fruit flies (allover the world) were controlled with the chemical pesticides that caused many healthy, environmentally and economically problems for man, natural enemies and fruit-producer countries. For this reasons many researchers used the plant extracts as naturally-safe materials for controlling insects (Rajendran & Gopalan, 1979, Smith & Secoy, 1981, Steffens & Schmutterer, 1982, Barakat *et al.*, 1985, Chan & Tam, 1985, Guirguis *et al.*, 1989, Kelany *et al.*, 1991, El-Abgy *et al.*, 1997, Pandey & Singh, 1998, Di Ilio *et al.*, 1999, El-Doksch & El-Sebae, 2005, El-Doksch & El-Sherif, 2005 and Fetoh *et al.*, 2005).

The aim of this work is to evaluate the insecticidal activity of certain plant extracts against both pupal and adult stages of the two species of tephritid fruit flies *C. capitata* and *B. zonata*.

MATERIALS AND METHODS

1- Insect Used

The two stages of pupae and adult used for treatments were obtained from cultures of the laboratory strains of both *C. capitata* and *B. zonata* that mass reared according to Awadallah and El-Hakim (1983) and Afia *et al.* (2005), respectively.

2- Plant Extracted

The insecticidal effect of aqueous dilutions of certain parts of six wild plant species (Table, 1) was tested against the two stages of the studied insects. The parts of the tested plants collected from the field of Horticulture Research Institute were air-dried under laboratory conditions of $25 \pm 3^\circ\text{C}$ and $70 \pm 5\%$ R.H. Afterthat, 20 gm of each were ground with ethanolic extraction (300 ml of commercial ethanol alcohol 75% for each plant) using Sockcehl unit.

Table 1. List of plant species under investigation.

	Scientific name	Family	Common name	Used part
1	<i>Ambrosia maritima</i>	Compositae	Absinthe	Flowers
2	<i>Allium sativum</i>	Liliaceae	Garlic	Bulbis
3	<i>Mentha spicata</i>	Labiatae	Spearmint	Leaves
4	<i>Myoporum pictum</i>	Myopaceae	Bazromia	Flowers
5	<i>Nerium oleander</i>	Apocynaceae	Nerium	Leaves
6	<i>Rosa gallica</i>	Rosaceae	Rose	Leaves

3- Procedure

Thirty individuals replicated three times (10 individuals/ replicate) of both pupal (3-day old) and adult (2-day after emergence) stages of the two tephritid species were sprayed with 0.5 ml of different aqueous concentrations of the tested extracts using an atomizer instrument. Three concentrations (5, 10 and 20%) were used for adults, sprayed in tubes (10 cm in length and 2 cm in diameter), whereas pupae were treated in Petri dishes (9 cm in diameter) with four concentrations of 25, 50, 75 and 100 %. The individuals of untreated control were sprayed with tap water. After 3 hours of treatment, pupae were put in clean Petri dishes to emergence, whereas adults, were transferred by an aspirator to clean small cages provided with sugar and water for feeding and examined after 24, 48 and 72 hours of treatment. For both pupae and adults, the dead individuals were recorded. Pupal mortality was corrected with Abbott's formula (1925). Both LC_{50} and LC_{90} values of the tested extracts were obtained from dosage-mortality regression lines drawn according to the method of Bliss (1938). The relative efficiency as toxicity index was calculated according to Sun

(1950). On the other hand, the cumulative mortality during 72 hours after treatments was calculated for adults. Analysis of variance was conducted to test significance between treatments using "F" test and L.S.D. values according to Snedecor (1957).

RESULTS AND DISCUSSION

1- Toxicity against Pupae

Data in Table (2) and Figs. (1 and 2) indicate the contact toxic effect of the six tested plants to 3- day old pupae of both *C. capitata* and *B. zonata*. Based on LC₅₀ and LC₉₀ values, leaves extract of *R. gallica* was the most potent against pupae of *C. capitata* recording the lowest values of 46 and 74%, respectively. Whereas leaves extract of *M. pictum* was the least effective showing LC₅₀ and LC₉₀ values of 81 and 88%, consecutively. The other extracts had intermediate values of LC₅₀ and LC₉₀ that ranged between 52- 81 and 77- 88, successively. The slope values of the toxicity lines reveal that *M. pictum* had the steepest toxicity line and *M. spicata* had the flattest one. As shown in Table (2), the toxicity index of the tested plant extracts ranged between 56.79- 97.87% (at LC₅₀) and 84.09- 96.10% (at LC₉₀) from leaves extract of *R. gallica* that was considered the standard.

Respecting the toxic effect on *B. zonata*, the bulbs extract of *A. sativum* was the standard recording the lowest values of both LC₅₀ and LC₉₀ of 28 and 42, successively. The efficiency of the other tested extracts was descendingly arranged as *N. oleander*, *A. maritima*, *R. gallica*, *M. pictum* and *M. spicata*. The values of both LC₅₀ and LC₉₀ were 40, 62, 42, 68, 46, 70, 61, 74 and 78, 92, respectively. The corresponding values of toxicity index were 70.00, 67.74, 66.67, 61.76, 60.87, 60.00, 45.90, 56.76 and 35.90, 54.65% of the standard extract of *A. sativum*, consecutively. The slope values of toxicity lines of the six examined extracts ranged between 6.49- 18.18 indicated that *M. spicata* had the steepest toxicity line, but *A. maritima* had the flattest one.

Based on LC₅₀ and LC₉₀ values, *B. zonata* was more susceptible to *A. maritima*, *A. sativum*, *M. pictum* and *N. oleander* than *C. capitata*, but the reverse was true with *M. spicata*. The response of the two tephritid species to *R. gallica* was equal and nearly equal according to values of LC₅₀ and LC₉₀, consecutively. In this respect, Rajendran and Gopalan (1979) stated that *A. sativum* had insecticidal properties against larvae of *Spodoptera litura*.

Table 2. Toxicity of certain plant extracts against 3-day old pupae of both *ceratitis capitata* and *B. zonata*

Plant extract	<i>C. capitata</i>					<i>B. zonata</i>				
	LC ₅₀	LC ₉₀	Slope	Toxicity index at		LC ₅₀	LC ₉₀	Slope	Toxicity index at	
				LC ₅₀	LC ₉₀				LC ₅₀	LC ₉₀
<i>A. maritima</i>	52	78	6.95	88.46	94.87	42	68	6.49	66.67	61.76
<i>A. sativum</i>	79	88	33.97	58.23	84.09	28	42	7.23	100	100
<i>M. spicata</i>	47	80	5.94	97.87	92.50	78	92	18.18	35.90	45.65
<i>M. pictum</i>	81	88	36.51	56.79	84.09	61	74	14.68	45.90	56.76
<i>N. oleander</i>	62	77	13.18	74.19	96.10	40	62	6.64	70.00	67.74
<i>R. gallica</i>	46	74	6.28	100	100	46	70	7.11	60.87	60.00

2- Cumulative Mortality of Adults

The cumulative mortality of 2-day old adults of both *C. capitata* and *B. zonata* sprayed with the examined plant extracts was shown in Table (3). Statistical analysis of variance of the obtained data revealed that there were high significant differences between the treated and untreated individuals. In case of *C. capitata* the differences between the six tested extracts were insignificant, but in case of *B. zonata*, the extract of *A. maritima* significantly varied with the others. Leaves extract of *N. oleander* was the most efficient to newly emerged adults of both *C. capitata* and *B. zonata* showing the highest values of cumulative mortality (after 72 hours) of 55.93 and 90.00%, whereas flowers extract of *A. maritima* recorded the lowest % cumulative mortalities of 45.19 and 57.41, respectively. The other extracts showed intermediate values of cumulative mortality that ranged between 46.29- 49.26% (in case of *C. capitata*) and 79.63- 87.41% (in case of *B. zonata*). Data in Table (3) show that adults of *C. capitata* were more tolerant to the tested extracts than that of *B. zonata*, where the grand averages of cumulative mortality for the six tested extracts were lower with *C. capitata* than that recorded with *B. zonata*. In this respect Barakat *et al.* (1985) reported that the acetone extract of black pepper was the most effective against *C. capitata* adults. Recently, Fetoh *et al.* (2005) evaluated different concentrations of ethanolic- leaves extract of *Calotropis procera* against adults of *Dacus ciliatus* and *B. zonata*. They stated that the extract was effective against the two species and *B. zonata* was more susceptible than *D. ciliatus*.




Fig. 1. LC-P Lines of six plant extracts against 3-day pupae of *Ceratitis Capitata*.




Fig. 2. LC-P Lines of six plant extracts against 3-day pupae of *Bactrocera Zonata*.

Table 3. Cumulative mortality of 2-day old adults of both *Ceratitis capitata* and *Bactrocera zonata* sprayed with certain plant extracts.

Plant extract	Conc. %	% Mortality after (hour)									
		<i>C. capitata</i>					<i>B. zonata</i>				
		24	48	72	Aver.	Grand aver.	24	48	72	Aver.	Grand aver.
<i>A. maritima</i>	5	13.33	46.67	83.33	47.78		10.00	40.00	96.67	48.89	
	10	30.00	46.67	66.67	47.78	45.19A	30.00	50.00	96.67	58.89	57.41B
	20	6.67	50.00	63.33	40.00		36.67	60.00	96.67	64.45	
<i>A. sativum</i>	5	10.00	50.00	63.33	41.11		30.00	60.00	96.67	62.22	
	10	10.00	50.00	70.00	43.33	46.29A	70.00	80.00	96.67	82.22	79.63A
	20	23.33	60.00	80.00	54.44		90.00	96.67	96.67	94.45	
<i>M. spicata</i>	5	16.67	56.67	73.33	48.89		40.00	90.00	96.67	75.56	
	10	20.00	56.67	73.33	50.00	47.78A	70.00	90.00	96.67	85.56	82.97A
	20	20.00	43.33	70.00	44.44		70.00	96.67	96.67	87.78	
<i>M. pictum</i>	5	13.33	56.67	73.33	47.78		63.33	76.67	93.33	77.78	
	10	40.00	50.00	86.67	58.89	49.26A	60.00	80.00	96.67	78.89	82.59A
	20	20.00	40.00	63.33	41.11		80.00	96.67	96.67	91.11	
<i>N. oleander</i>	5	13.33	63.33	70.00	48.89		66.67	90.00	96.67	84.45	
	10	43.33	63.33	80.00	62.22	55.93A	80.00	90.00	96.67	88.89	90.00A
	20	43.33	60.00	66.67	56.67		96.67	96.67	96.67	96.67	
<i>R. gallica</i>	5	16.67	36.67	76.67	43.34		70.00	93.33	96.67	86.67	
	10	33.33	53.33	93.33	60.00	49.26A	70.00	93.33	96.67	86.67	87.41A
	20	20.00	43.33	70.00	44.44		80.00	90.00	96.67	88.89	
Control	0	0	0	0	0	0B	0	0	0	0	0C
F. test						**					**

REFERENCES

- 1- Abbott, W.S. 1925. A method of computing the effectiveness of an insecticide. J. Econ. Entomol., 18: 265- 267.
- 2- Afia, Y.I., M.A. El-Sheikh, A.G. Hashem and S. El-Nagar. 2005. Influence of different constant temperatures on the biological aspects of the two fruit flies (Diptera: Tephritidae) in Egypt. 1. On survival and developmental rates of immature stages. Egypt. J. Appl. Sci., 20 (7): 257- 276.
- 3- Awadallah, A.M. and Aida M. El-Hakim. 1987. Methods for mass production of the Mediterranean fruit fly *Ceratitis capitata* Wied. II-Methods for producing adults. Zagazig J. Agric. Res., 14 (1): 709- 719.
- 4- Barakat, A.A., H.S.M. Fahmy, M.A. Kandil and N.M.M. Ebrahim. 1985. Toxicity of the extracts of black-pepper, cumin, fennel, chamomille and lupin against *Drosophila melanogaster*, *Ceratitis capitata* and *Spodoptera littoralis*. Indian J. Agric. Sci., 55 (2): 116- 120.
- 5- Bliss, C.I. 1938. The determination of dosage mortality curves from small numbers. Quart. J. Phar. and Pharmacology, 11(2): 192.
- 6- Chan, H.T.Jr. and S.Y.T. Tam. 1985. Toxicity of α - tomatine to larvae of the Mediterranean fruit fly (Diptera: Tephritidae). J. Econ. Entomol., 78: 305- 307.
- 7- Di Ilio, Vincenzo, Massimo Cristofaro, Daniela Marchini, Paola Nobili and Romano Dallai. 1999. Effects of a neem compound on the fecundity and longevity of *Ceratitis capitata* (Diptera: Tephritidae). J. Econ. Entomol., 92 (1): 76- 82.
- 8- El-Abgy, Elham, A.A. El-Fishawi and R.G. Abou-El-Ela. 1997. Antagonistic actions of *Ageratum conyzoides* on *Culex pipiens pipiens* in Egypt. Egypt. J. Appl. Sci., 12 (1): 290- 305.
- 9- El-Doksch, H.A. and A.H. El-Sebae. 2005. Plant natural products as a source of new and environmentally safe pesticides within IPM programmes. Egypt. J. Agric. Res., 83 (3): 1127- 1145.
- 10- El-Doksch, H.A. and M.G. El-Sherif. 2005. Inhibition of growth and development of cotton leaf worm (Lepidoptera: Noctuidae) larvae by β - Carboline alkaloid. J. Agric. Sci. Mansoura Univ., 30 (1): 401- 408.
- 11- Fetoh, Badr El-Sabah A., Gehad M. Mousa and Tahany G. Mohamaden. 2005. Utilization of El-Oshar plant extract as a novel and safe approach in controlling

- the pumpkin fly, *Dacus ciliatus* Loew and the peach fruit fly, *Bactrocera zonata* Saunders (Diptera: Tephritidae) in Egypt. *Egypt. J. Agric. Res.*, 83 (1): 285-291.
- 12- Guirguis, M.W., K.M. Gouhar, W.M. Watson and R.M. Salem. 1989. Toxicity and antifeedant activity of wild plant extracts against some insect pests. *Agric. Res. Rev.*, 67 (1): 1-12.
 - 13- Kelany, I.M., M.A. El-Deeb and A.A. El-Fishawi. 1991. The effect of aqueous neem, *Azadirachta indica* A. Juss, seed kernel extract on the house fly, *Musca domestica* L. *Egypt. J. Biol. P. Cont.*, 1 (1): 121 -128.
 - 14- Pandey, K.P. and Suchita Singh. 1998. Evaluation of natural plant product as an insecticide against top borer *Scirpophaga excerptalis* Wlk. in sugarcane. *Entomon*, 23 (3): 235- 237.
 - 15- Rajendran, B. and M. Gopalan. 1979. Note on the insecticidal properties of certain plant extracts. *Indian J. Agric. Sci.*, 49 (4) 295- 297.
 - 16- Smith, A. and M. Secoy. 1981. Plants used for agricultural pest control in western Europe before 1850. *Chemistry & Industry*, No. 1: 12- 17.
 - 17- Snedecor, G.W. 1957. *Statistical methods applied to experiments in agriculture and biology*. The Iowa State Colledge Press, Amer., Iowa, 5th ed.
 - 18- Steffens, R.J. and H. Schmutterer. 1982. The effect of a crude methanolic neem (*Azadirachta indica*) seed kernel extract on metamorphosis and quality of adults of the med fly (*Ceratitis capitata*) (Wied.). *Z. Angew. Entomol.*, 94: 98- 103.
 - 19- Sun, Y.P. 1950. Toxicity index- An improved method of comparing the relative toxicity of insecticides. *J. Econ. Entomol.*, 43 (1): 45- 53.

كفاءة بعض المستخلصات النباتية ضد ذبابة فاكهة البحر المتوسط وذبابة ثمار الخوخ

أحمد حسين حنفي^١، أناس عبد العزيز أحمد^١، أحمد محمود زكي مسلم^٢،
عبد الحميد محمد إبراهيم^١

١. معهد بحوث وقاية النباتات- الصباحية- الإسكندرية

٢. معهد بحوث وقاية النباتات- مركز البحوث الزراعية

تم اختبار تأثير مستخلصات ٦ نباتات برية هي البازروميا- الثوم- النعناع البلدي-
الدمسيصة- النغلة- الورد البلدي الأحمر كمواد سامة بالملامسة ضد العذارى عمر ٣ أيام والأطوار
الكاملة لحشري ذبابة فاكهة البحر المتوسط وذبابة ثمار الخوخ. كانت مستخلصات أوراق الورد
الأحمر وفصوص الثوم الأكثر فعالية ضد عذارى ذبابة فاكهة البحر المتوسط وذبابة ثمار الخوخ علي
الترتيب. ومن جهة أخرى كان مستخلص أوراق النغلة الأكثر كفاءة ضد الحشرات الكاملة لذبابتي
فاكهة البحر المتوسط وثمار الخوخ حيث سجل أعلى قيمة للموت التراكمي (بعد ٧٢ ساعة) وهي
٥٥,٩٢%، ٩٠% علي التوالي. وكانت الحشرات الكاملة لذبابة الخوخ أكثر حساسية للمستخلصات
المختبرة منها لذبابة فاكهة البحر المتوسط.